The Journal of Head Trauma Rehabilitation®

Cumulative Contents to Volume 19

19:1 January–February 2004
Pain Management
Nathan D. Zasler, MD, FAADEP, FAAPM&R, DAAPM, CIME, and Michael F. Martelli, PhD, DAAPM
Issue Editors

19:2 March–April 2004
Spasticity and the Upper Motor Neuron Syndrome in Traumatic Brain Injury
Elie P. Elovic, MD, and Mel B. Glenn, MD
Issue Editors

19:3 May–June 2004
Focus on Clinical Research and Practice, Part 1
Bruce Caplan, PhD, ABPP, and Ross D. Zafonte, DO
Issue Editors

19:4 July–August 2004
Long-term Neurologic Complications of Traumatic Brain Injuries
Ramon Diaz-Arrastia, MD, PhD
Issue Editor

19:5 September–October 2004
Focus on Clinical Research and Practice, Part 2
Bruce Caplan, PhD, ABPP
Issue Editor

19:6 November–December 2004
Social Participation After Traumatic Brain Injury
Keith D. Cicerone, PhD, ABPP
Issue Editor

Author Index

A

Addis PE: *See* Cahill LM

Agostini M: *See* Hudak AM

Arndt S: *See* Jorge R

Ashman T: *See* Brown M

B

Barisa MT: *Traumatic Brain Injury: Methods for Clinical and Forensic Neuropsychiatric Assessment*, by Granacher RP (Book Review), 19(3):284

Bell KR, Hoffman JM, Doctor JN, Powell JM, Esselman P, Bombardier C, Fraser R, Dikmen S: Development of a Telephone Follow-up Program for Individuals Following Traumatic Brain Injury (Edited by Zafonte R), 19(6):502

Bender MC: *See* Martelli MF

Bertrand R: *See* Dumont C

Bogner J: *See* Corrigan JD

Bombardier C: *See* Bell KR

Booker K: *See* Hudak AM

Branca B, Lake III AE: Psychological and Neuropsychological Integration in Multidisciplinary Pain Management After TBI, 19(1):40

Brener I, Harman JS, Kelleher KJ, Yeates KO: Medical Costs of Mild to Moderate Traumatic Brain Injury in Children, 19(5):405

Brown M, Dijkers MPJM, Gordon WA, Ashman T, Charatz H, Cheng Z: Participation Objective, Participation Subjective: A Measure of Participation Combining Outsider and Insider Perspectives, 19(6):459

Brunner RC: *See* Meythaler JM

Buchhofer R: *See* Hart T

Burrington B: *See* Fann, J

Bushnik T, Englander J, Duong T: Medical and Social Issues Related to Posttraumatic Seizures in Persons With Traumatic Brain Injury, 19(4):296

Bushnik T: *See* Hammond FM

Bérubé JE: The Annual Congressional Brain Injury Awareness Fair (Edited by Bérubé JE), 19(1):82

C

Caesar RR: *See* Hudak AM

Cahill LM, Turner AB, Stabler PA, Addis PE, Theodoros DG, Murdoch BE: An Evaluation of Continuous Positive Airway Pressure (CPAP) Therapy in the Treatment of Hypernasality Following Traumatic Brain Injury: A Report of 3 Cases, 19(3):241

Carnevale GJ: *Neuropsychosocial Intervention: The Practical Treatment of Severe Behavioral Dyscontrol after Acquired Brain Injury*, by Karol RL (Book Review), 19(2):188

Charatz H: *See* Brown M

Cheng Z: *See* Brown M

Cicerone KD: Participation as an Outcome of Traumatic Brain Injury Rehabilitation, 19(6):494

Cicerone KD: Social Participation After Traumatic Brain Injury (Preface), 19(6):429

Clayton W: *See* Meythaler JM

Corrigan JD, Bogner J: Latent Factors in Measures of Rehabilitation Outcomes After Traumatic Brain Injury, 19(6):445

Corrigan JD, Whiteneck G, Mellick D: Perceived Needs Following Traumatic Brain Injury, 19(3):205

Corrigan JD: *See* Hammond FM

Crespo-Fecarro B: *See* Jorge R

Cusick CP: *See* Whiteneck GG

D

Davis LK: *See* Meythaler JM

Dewey Jr RB: *See* O'Suilleabhain P

Diaz-Arrastia R: Long-term Neurologic Complications of Traumatic Brain Injuries (Preface), 19(4):289

Diaz-Arrastia R: *See* Hudak AM

Dijkers MPJM: *See* Brown M

Dikmen S: *See* Bell KR

Doctor JN: *See* Bell KR

Dumont C, Gervais M, Fougeyrollas P, Bertrand R: Toward an Explanatory Model of Social Participation for Adults With Traumatic Brain Injury, 19(6):431

Duong T: *See* Bushnik T

E

Elovic EP, Glenn MB: Spasticity and the Upper Motor Neuron Syndrome in Traumatic Brain Injury (Preface), 19(2):87

Elovic EP, Simone LK, Zafonte R: Outcome Assessment for Spasticity Management in the Patient With Traumatic Brain Injury: The State of the Art, 19(2):155

Elovic EP: Anterior Pituitary Dysfunction After Traumatic Brain Injury, Part II (Edited by Glenn MB), 19(2):184

Elovic EP: *See* Zafonte R

Englander J: *See* Bushnik T

Esquenazi A: Evaluation and Management of Spastic Gait in Patients With Traumatic Brain Injury, 19(2):109

Esselman P: *See* Bell KR

F

Fann J, Burrington B, Leonetti A, Jaffe K, Katon W, Thompson R: Abstract of psychiatric illness following traumatic brain injury in an adult health maintenance organization population (Abstract by Laborde A), 19(4):349

Fichtenberg NL: *See* Mahmood O

Forget R: *See* Gagnon I

Fougeyrollas P: *See* Dumont C

Fraser R: *See* Bell KR

Friedman D: *See* Gagnon I

G

Gagnon I, Swaine B, Friedman D, Forget R: Visuomotor Response Time in Children With a Mild Traumatic Brain Injury, 19(5):391

Gerhart KA: *See* Whiteneck GG

Gervais M: *See* Dumont C

Giacino JT, Trott CT: Rehabilitative Management of Patients With Disorders of Consciousness: Grand Rounds, 19(3):254

Glenn MB, Goldstein R, Selleck EA, Rotman M: Characteristics of Facility-based Community Integration Programs for People With Brain Injury, 19(6):482

Glenn MB: *See* Elovic EP

Goldstein R: *See* Glenn MB

Gordon WA: *See* Brown M

Greenspan AI: *See* Phillips VL

Guin-Renfroe S: *See* Meythaler JM

H

Hammond FM, Hart T, Bushnik T, Corrigan JD, Sasser H: Change and Predictors of Change in Communication, Cognition, and Social Function Between 1 and 5 Years After Traumatic Brain Injury, 19(4):314

Hanks RA: *See* Mahmood O

Harman JS: *See* Brener I

Harper CR: *See* Hudak AM

Hart T, Buchhofer R, Vaccaro M: Portable Electronic Devices as Memory and Organizational Aids After Traumatic Brain Injury: A Consumer Survey Study, 19(5):351

Hart T: *See* Hammond FM

Hartman ET: *See* Ivanhoe CB

Hecht JS: Occipital Nerve Blocks in Postconcussive Headaches: A Retrospective Review and Report of Ten Patients, 19(1):58

Hoffman JM: *See* Bell KR

Hudak AM, Trivedi K, Harper CR, Booker K, Caesar RR, Agostini M, Van Ness PC, Diaz-Arrastia R: Evaluation of Seizure-like Episodes in Survivors of Moderate and Severe Traumatic Brain Injury, 19(4):290

Hunt E: *See* Powell JM

I

Ivanhoe CB, Hartman ET: Clinical Caveats on Medical Assessment and Treatment of Pain After TBI, 19(1):29

J

Jaffe K: *See* Fann, J

Jorge R, Robinson R, Crespo-Fecarro B, Arndt S: Major depression following traumatic brain injury (Abstract by Laborde A), 19(4):349

K

Katon W: *See* Fann, J

Keenan MAE, Mehta S: Neuro-Orthopedic Management of Shoulder Deformity and Dysfunction in Brain-Injured Patients: A Novel Approach, 19(2):143

Kelleher KJ: *See* Brener I

Kirsch NL, Shenton M, Spirl E, Simpson R, LoPresti E, Schreckenghost D: An Assistive-Technology Intervention for Verbose Speech After Traumatic Brain Injury: A Single Case Study, 19(5):366

L

Lake III AE: *See* Branca B

Lehtonen S: *See* Phillips VL

Leonetti A: *See* Fann, J

Lombard L: *See* Zafonte R

Long E: *See* Simpson G

LoPresti E: *See* Kirsch NL

Lynch WJ: A New Neuropsychological Test Battery: The NAB (Edited by Lynch WJ), 19(2):180

Lynch WJ: Determination of Effort Level, Exaggeration, and Malingering in Neurocognitive Assessment (Edited by Lynch WJ), 19(3):277

M

Mahmood O, Rapport LJ, Hanks RA, Fichtenberg NL: Neuropsychological Performance and Sleep Disturbance Following Traumatic Brain Injury, 19(5):378

Martelli MF, Zasler ND, Bender MC, Nicholson K: Psychological, Neuropsychological, and Medical Considerations in Assessment and Management of Pain, 19(1):10

Martelli MF: *See* Nicholson K

Martelli MF: *See* Zasler ND

Mayer NH: Choosing Upper Limb Muscles for Focal Intervention After Traumatic Brain Injury, 19(2):119

Mehta S: *See* Keenan MAE

Mellick D: *See* Corrigan JD

Meythaler JM, Clayton W, Davis LK, Guin-Renfroe S, Brunner RC: Orally Delivered Baclofen to Control Spastic Hypertonia in Acquired Brain Injury, 19(2):101

Morgan A, Ward E, Murdoch B: Clinical Characteristics of Acute Dysphagia in Pediatric Patients Following Traumatic Brain Injury, 19(3):226

Moseley AM, Lanzarone S, Bosman JM, van Loo MA, de Bie RA, Hassett L: Ecological Validity of Walking Speed Assessment After Traumatic Brain Injury: A Pilot Study (Edited by Caplan B), 19(4):341

Murdoch B: *See* Morgan A

Murdoch BE: *See* Cahill LM

N

Nicholson K, Martelli MF: The Problem of Pain, 19(1):2

Nicholson K: *See* Martelli MF

O

O'Suilleabhain P, Dewey Jr RB: Movement Disorders After Head Injury: Diagnosis and Management, 19(4):305

P

Pepping M: *See* Powell JM

Perna R: Benzodiazepines and Antipsychotics: Cognitive Side Effects (Edited by Glenn MB), 19(6):516

Phillips VL, Greenspan AI, Stringer AY, Stroble AK, Lehtonen S: Severity of Injury and Service Utilization Following Traumatic Brain Injury: The First 3 Months, 19(3):217

Powell JM, Hunt E, Pepping M: Collaboration Between Cognitive Science and Cognitive Rehabilitation: A Call for Action, 19(3):266

Powell JM: *See* Bell KR

R

Rapport LJ: *See* Mahmood O

Robinson R: *See* Jorge R

Rotman M: *See* Glenn MB

S

Sasser H: *See* Hammond FM

Schreckenghost D: *See* Kirsch NL

Selleck EA: *See* Glenn MB

Shenton M: *See* Kirsch NL

Simone LK: *See* Elovic EP

Simpson G, Long E: An Evaluation of Sex Education and Information Resources and Their Provision to Adults With Traumatic Brain Injury, 19(5):413

Simpson R: *See* Kirsch NL

Spirl E: *See* Kirsch NL

Stabler PA: *See* Cahill LM

Stringer AY: *See* Phillips VL

Stroble AK: *See* Phillips VL

Stuart M: Fighting the Silent Epidemic: The Florida Brain and Spinal Cord Injury Program (Edited by Zafonte R), 19(4):329

Swaine B: *See* Gagnon I

T

Theodoros DG: *See* Cahill LM

Thompson R: *See* Fann, J

Tovino SA: *See* Winslade WJ

Trivedi K: *See* Hudak AM

Trott CT: *See* Giacino JT

Turner AB: *See* Cahill LM

V

Vaccaro M: *See* Hart T

Van Ness PC: *See* Hudak AM

W

Walker WC: Pain Pathoetiology After TBI: Neural and Nonneural Mechanisms, 19(1):72

Ward E: *See* Morgan A

Whiteneck G: *See* Corrigan JD

Whiteneck GG, Gerhart KA, Cusick CP: Identifying Environmental Factors That Influence the Outcomes of People With Traumatic Brain Injury, 19(3):191

Winslade WJ, Tovino SA: Research With Brain-Injured Subjects (Edited by Winslade WJ), 19(6):513

Winslade WJ: Research on Minimally Conscious Patients: Innovation or Exploitation? (Edited by Winslade WJ), 19(2):178

Y

Yeates KO: *See* Brener I

Z

Zafonte R, Elovic EP, Lombard L: Acute Care Management of Post-TBI Spasticity, 19(2):89

Zafonte R: *See* Elovic EP

Zasler ND, Martelli MF: Pain Management (Preface), 19(1):1

Zasler ND: *See* Martelli MF

Zasler ND: *See* Martelli MF

Zimmer C: The vegetative state: guidance on diagnosis and management (Abstract by Laborde A), 19(1):85

Zimmer C: What if there is something going on in there (Abstract by Laborde A), 19(1):85

Subject Index

A

Abstracts

Abstract of psychiatric illness following traumatic brain injury in an adult health maintenance organization population, Fann J, Burrington B, Leonetti A, et al (abstracted by Laborde A), 19(4):349

Major depression following traumatic brain injury, Jorge R, Robinson R, Crespo-Fecarro B, Arndt S (abstracted by Herman CD), 19(4):349

The vegetative state: guidance on diagnosis and management (abstracted by Laborde A), 19(1):85

What if there is something going on in there, Zimmer C (abstracted by Laborde A), 19(1):85

Acute care

Acute care management of post-TBI spasticity, Zafonte R, Elovic EP, Lombard L, 19(2):89–100

Acute pain

Pain pathoetiology after TBI: neural and nonneural mechanisms, Walker WC, 19(1):72–81

Adhesive capsulitis

Neuro-orthopedic management of shoulder deformity and dysfunction in brain-injured patients: a novel approach, Keenan MAE, Mehta S, 19(2):143–154

Adrenocorticotropic hormone (ACTH) deficiency

Anterior pituitary dysfunction after traumatic brain injury: Part II, Elovic EP, 19(2):184–187

Anoxia

Orally delivered baclofen to control spastic hypertonia in acquired brain injury, Meythaler JM, Clayton W, Davis LK, Guin-Renfroe S, Brunner RC, 19(2):101–108

Antipsychotic drugs

Benzodiazepines and antipsychotics: cognitive side effects, Perna R, 19(6):516–518

Arms

Choosing upper limb muscles for focal intervention after traumatic brain injury, Mayer NH, 19(2):119–142

Ashworth Scale

Outcome assessment for spasticity management in the patient with traumatic brain injury: the state of the art, Elovic EP, Simone LK, Zafonte R, 19(2):155–177

Assistive technology

An assistive technology intervention for verbose speech after traumatic brain injury: a single case study, Kirsch NL, Shenton M, Spirl E, et al, 19(5):366–377

Portable electronic devices as memory and organizational aids after traumatic brain injury: a consumer survey study, Hart T, Buchhofer R, Vaccaro M, 19(5):351–365

B

Baclofen

Acute care management of post-TBI spasticity, Zafonte R, Elovic EP, Lombard L, 19(2):89–100

Clinical caveats on medical assessment and treatment of pain after TBI, Ivanhoe CB, Hartman ET, 19(1):29–39

Orally delivered baclofen to control spastic hypertonia in acquired brain injury, Meythaler JM, Clayton W, Davis LK, Guin-Renfroe S, Brunner RC, 19(2):101–108

Balance

Outcome assessment for spasticity management in the patient with traumatic brain injury: the state of the art, Elovic EP, Simone LK, Zafonte R, 19(2):155–177

Behavioral control

Perceived needs following traumatic brain injury, Corrigan JD, Whiteneck G, Mellick D, 19(3):205–216

Benzodiazepines

Benzodiazepines and antipsychotics: cognitive side effects, Perna R, 19(6):516–518

Biopsychosocial assessment

Psychological, neuropsychological, and medical considerations in assessment and management of pain, Martelli MF, Zasler ND, Bender MC, Nicholson K, 19(1):10–28

Book reviews

Neuropsychosocial Intervention: The Practical Treatment of Severe Behavioral Dyscontrol After Acquired Brain Injury: Karol RL (reviewed by Carnevale GJ), 19(2):188–189

Traumatic Brain Injury: Methods for Clinical and Forensic Neuropsychiatric Assessment: Granacher RP (reviewed by Barisa MT), 19(3):284–285

Botulinum toxin (Botox)

Clinical caveats on medical assessment and treatment of pain after TBI, Ivanhoe CB, Hartman ET, 19(1):29–39

Brain injury. *See also* Pediatric brain injury

Acute care management of post-TBI spasticity, Zafonte R, Elovic EP, Lombard L, 19(2):89–100

An assistive technology intervention for verbose speech after traumatic brain injury: a single case study, Kirsch NL, Shenton M, Spirl E, et al, 19(5):366–377

An evaluation of continuous positive airway pressure (CPAP) therapy in the treatment of hypernasality following traumatic brain injury: a report of 3 cases, Cahill LM, Turner AB, Stabler PA, et al, 19(3):241–253

An evaluation of sex education and information resources and their provision to adults with

traumatic brain injury, Simpson G, Long E, 19(5):413–428

The Annual Congressional Brain Injury Awareness Fair, Bérubé JE, 19(1):82–84

Anterior pituitary dysfunction after traumatic brain injury: Part II, Elovic EP, 19(2):184–187

Benzodiazepines and antipsychotics: cognitive side effects, Perna R, 19(6):516–518

Change and predictors of change in communication, cognition, and social function between 1 and 5 years after traumatic brain injury, Hammond FM, Hart T, Bushnik T, et al, 19(4):314–328

Characteristics of facility-based community integration programs for people with brain injury, Glenn MR, Goldstein R, Selleck EA, Rotman M, 19(6):482–493

Choosing upper limb muscles for focal intervention after traumatic brain injury, Mayer NH, 19(2):119–142

Clinical caveats on medical assessment and treatment of pain after TBI, Ivanhoe CB, Hartman ET, 19(1):29–39

Collaboration between cognitive science and cognitive rehabilitation: a call for action, Powell JM, Hunt E, Pepping M, 19(3):266–276

Determination of effort level, exaggeration, and malingering in neurocognitive assessment, Lynch WJ, 19(3):277–283

Development of a telephone follow-up program for individuals following traumatic brain injury, Bell KR, Hoffman JM, Doctor JN, et al, 19(6):502–512

Ecological validity of walking speed assessment after traumatic brain injury: a pilot study, Moseley AM, Lanzarone S, Bosman JM, et al, 19(4):341–348

Evaluation and management of spastic gait in patients with traumatic brain injury, Esquenazi A, 19(2):109–118

Evaluation of seizure-like episodes in survivors of moderate and severe traumatic brain injury, Hudak AM, Trivedi K, Harper CR, et al, 19(4):290–295

Fighting the silent epidemic: the Florida Brain and Spinal Cord Injury Program, Stuart M, 19(4):329–340

Identifying environmental factors that influence the outcomes of people with traumatic brain injury, Whiteneck GG, Gerhart KA, Cusick CP, 19(3):191–204

Latent factors in measures of rehabilitation outcomes after traumatic brain injury, Corrigan JD, Bogner J, 19(6):445–458

Medical and social issues related to posttraumatic seizures in persons with traumatic brain injury, Bushnik T, Englander J, Duong T, 19(4):296–304

Neuro-orthopedic management of shoulder deformity and dysfunction in brain-injured patients: a novel approach, Keenan MAE, Mehta S, 19(2):143–154

Neuropsychological performance and sleep disturbance following traumatic brain injury, Mahmood O, Rapport LJ, Hanks RA, Fichtenberg NL, 19(5):378–390

Occipital nerve blocks in postconcussive headaches: a retrospective review and report of ten patients, Hecht JS, 19(1):58–71

Orally delivered baclofen to control spastic hypertonia in acquired brain injury, Meythaler JM, Clayton W, Davis LK, Guin-Renfroe S, Brunner RC, 19(2):101–108

Outcome assessment for spasticity management in the patient with traumatic brain injury: the state of the art, Elovic EP, Simone LK, Zafonte R, 19(2):155–177

Pain pathoetiology after TBI: neural and nonneural mechanisms, Walker WC, 19(1):72–81

Participation as an outcome of traumatic brain injury rehabilitation, Cicerone KD, 19(6):494–501

Perceived needs following traumatic brain injury, Corrigan JD, Whiteneck G, Mellick D, 19(3):205–216

Portable electronic devices as memory and organizational aids after traumatic brain injury: a consumer survey study, Hart T, Buchhofer R, Vaccaro M, 19(5):351–365

The problem of pain, Nicholson K, Martelli MF, 19(1):2–9

Psychological, neuropsychological, and medical considerations in assessment and management of pain, Martelli MF, Zasler ND, Bender MC, Nicholson K, 19(1):10–28

Psychological and neuropsychological integration in multidisciplinary pain management after TBI, Branca B, Lake AE III, 19(1):40–57

Research with brain-injured subjects, Tovino SA, 19(6):513–515

Severity of injury and service utilization following traumatic brain injury: the first 3 months, Phillips VL, Greenspan AI, Stringer AY, Stroble AK, Lehtonen S, 19(3):217–225

Toward an explanatory model of social participation for adults with traumatic brain injury, Dumont C, Gervais M, Fougeyrollas P, Bertrand R, 19(6):431–444

C

Case management

Fighting the silent epidemic: the Florida Brain and Spinal Cord Injury Program, Stuart M, 19(4):329–340

Cervicogenic headache

Occipital nerve blocks in postconcussive headaches: a retrospective review and report of ten patients, Hecht JS, 19(1):58–71

Chemodenervation

Choosing upper limb muscles for focal intervention after traumatic brain injury, Mayer NH, 19(2):119–142

Chorea

Movement disorders after head injury, O'Suilleabhain P, Dewey RB, Jr., 19(4):305–313

Chronic care

Fighting the silent epidemic: the Florida Brain and Spinal Cord Injury Program, Stuart M, 19(4):329–340

Chronic pain

Pain pathoetiology after TBI: neural and nonneural mechanisms, Walker WC, 19(1):72–81

The problem of pain, Nicholson K, Martelli MF, 19(1):2–9

Psychological and neuropsychological integration in multidisciplinary pain management after TBI, Branca B, Lake AE III, 19(1):40–57

Cognition

Change and predictors of change in communication, cognition, and social function between 1 and 5 years after traumatic brain injury, Hammond FM, Hart T, Bushnik T, et al, 19(4):314–328

Collaboration between cognitive science and cognitive rehabilitation: a call for action, Powell JM, Hunt E, Pepping M, 19(3):266–276

Cognitive compensation

Portable electronic devices as memory and organizational aids after traumatic brain injury: a consumer survey study, Hart T, Buchhofer R, Vaccaro M, 19(5):351–365

Cognitive disorders

An assistive technology intervention for verbose speech after traumatic brain injury: a single case study, Kirsch NL, Shenton M, Spirl E, et al, 19(5):366–377

Collaboration between cognitive science and cognitive rehabilitation: a call for action, Powell JM, Hunt E, Pepping M, 19(3):266–276

Cognitive impairment

Benzodiazepines and antipsychotics: cognitive side effects, Perna R, 19(6):516–518

Cognitive problems

Perceived needs following traumatic brain injury, Corrigan JD, Whiteneck G, Mellick D, 19(3):205–216

Collaboration

Collaboration between cognitive science and cognitive rehabilitation: a call for action, Powell JM, Hunt E, Pepping M, 19(3):266–276

Coma

Rehabilitative management of patients with disorders of consciousness: grand rounds, Giacino JT, Trott CT, 19(3):254–265

Communication

Change and predictors of change in communication, cognition, and social function between 1 and 5 years after traumatic brain injury, Hammond FM, Hart T, Bushnik T, et al, 19(4):314–328

Collaboration between cognitive science and cognitive rehabilitation: a call for action, Powell JM, Hunt E, Pepping M, 19(3):266–276

Development of a telephone follow-up program for individuals following traumatic brain injury, Bell KR, Hoffman JM, Doctor JN, et al, 19(6):502–512

Community integration

Characteristics of facility-based community integration programs for people with brain injury, Glenn MR, Goldstein R, Selleck EA, Rotman M, 19(6):482–493

Fighting the silent epidemic: the Florida Brain and Spinal Cord Injury Program, Stuart M, 19(4):329–340

Participation as an outcome of traumatic brain injury rehabilitation, Cicerone KD, 19(6):494–501

Participation objective, participation subjective: a measure of participation combining outsider and insider perspectives, Brown M, Dijkers PJM, Gordon WA, et al, 19(6):459–481

Concussion

Occipital nerve blocks in postconcussive headaches: a retrospective review and report of ten patients, Hecht JS, 19(1):58–71

Psychological and neuropsychological integration in multidisciplinary pain management after TBI, Branca B, Lake AE III, 19(1):40–57

Consciousness

Rehabilitative management of patients with disorders of consciousness: grand rounds, Giacino JT, Trott CT, 19(3):254–265

Research on minimally conscious patients: innovation or exploitation? Winslade WJ, 19(2):178–179

Consumer survey

Portable electronic devices as memory and organizational aids after traumatic brain injury: a consumer survey study, Hart T, Buchhofer R, Vaccaro M, 19(5):351–365

Continuous positive airway pressure (CPAP)

An evaluation of continuous positive airway pressure (CPAP) therapy in the treatment of hypernasality following traumatic brain injury: a report of 3 cases, Cahill LM, Turner AB, Stabler PA, et al, 19(3):241–253

Contracture

Neuro-orthopedic management of shoulder deformity and dysfunction in brain-injured patients: a novel approach, Keenan MAE, Mehta S, 19(2):143–154

D

Dantrolene sodium

Acute care management of post-TBI spasticity, Zafonte R, Elovic EP, Lombard L, 19(2):89–100

Deep venous thrombosis

Clinical caveats on medical assessment and treatment of pain after TBI, Ivanhoe CB, Hartman ET, 19(1):29–39

Determination (will)

Toward an explanatory model of social participation for adults with traumatic brain injury, Dumont C, Gervais M, Fougeyrollas P, Bertrand R, 19(6):431–444

Diazepam

Acute care management of post-TBI spasticity, Zafonte R, Elovic EP, Lombard L, 19(2):89–100

Disability

An evaluation of sex education and information resources and their provision to adults with traumatic brain injury, Simpson G, Long E, 19(5):413–428

Participation objective, participation subjective: a measure of participation combining outsider and insider perspectives, Brown M, Dijkers PJM, Gordon WA, et al, 19(6):459–481

Dynamism

Toward an explanatory model of social participation for adults with traumatic brain injury, Dumont C, Gervais M, Fougeyrollas P, Bertrand R, 19(6):431–444

Dysarthria

An evaluation of continuous positive airway pressure (CPAP) therapy in the treatment of hypernasality following traumatic brain injury: a report of 3 cases, Cahill LM, Turner AB, Stabler PA, et al, 19(3):241–253

Dysautonomia

Clinical caveats on medical assessment and treatment of pain after TBI, Ivanhoe CB, Hartman ET, 19(1):29–39

Dysphagia

Clinical characteristics of acute dysphagia in pediatric patients following traumatic brain injury, Morgan A, Ward E, Murdoch B, 19(3):226–240

Dystonia

Movement disorders after head injury, O'Suilleabhain P, Dewey RB, Jr., 19(4):305–313

E

Economic issues

Medical costs of mild to moderate traumatic brain injury in children, Brener I, Harman JS, Kelleher KJ, Yeates KO, 19(5):405–412

Effort level

Determination of effort level, exaggeration, and malingering in neurocognitive assessment, Lynch WJ, 19(3):277–283

Electromyography

Choosing upper limb muscles for focal intervention after traumatic brain injury, Mayer NH, 19(2):119–142

Employment

Perceived needs following traumatic brain injury, Corrigan JD, Whiteneck G, Mellick D, 19(3):205–216

Encephalopathy

Orally delivered baclofen to control spastic hypertonia in acquired brain injury, Meythaler JM, Clayton W, Davis LK, Guin-Renfroe S, Brunner RC, 19(2):101–108

Environmental barriers

Identifying environmental factors that influence the outcomes of people with traumatic brain injury, Whiteneck GG, Gerhart KA, Cusick CP, 19(3):191–204

Epilepsy

Evaluation of seizure-like episodes in survivors of moderate and severe traumatic brain injury, Hudak AM, Trivedi K, Harper CR, et al, 19(4):290–295

Medical and social issues related to posttraumatic seizures in persons with traumatic brain injury, Bushnik T, Englander J, Duong T, 19(4):296–304

Ethical issues

Research on minimally conscious patients: innovation or exploitation? Winslade WJ, 19(2):178–179

Research with brain-injured subjects, Tovino SA, 19(6):513–515

F

Fatigue

Toward an explanatory model of social participation for adults with traumatic brain injury, Dumont C, Gervais M, Fougeyrollas P, Bertrand R, 19(6):431–444

Financial burden

Medical costs of mild to moderate traumatic brain injury in children, Brener I, Harman JS, Kelleher KJ, Yeates KO, 19(5):405–412

Florida Brain and Spinal Cord Injury Program

Fighting the silent epidemic: the Florida Brain and Spinal Cord Injury Program, Stuart M, 19(4):329–340

Fractures

Clinical caveats on medical assessment and treatment of pain after TBI, Ivanhoe CB, Hartman ET, 19(1):29–39

Function

Outcome assessment for spasticity management in the patient with traumatic brain injury: the state of the art, Elovic EP, Simone LK, Zafonte R, 19(2):155–177

Functional assessment

Ecological validity of walking speed assessment after traumatic brain injury: a pilot study, Moseley AM, Lanzarone S, Bosman JM, et al, 19(4):341–348

Functional Independence Measure

Change and predictors of change in communication, cognition, and social function between 1 and 5 years after traumatic brain injury, Hammond FM, Hart T, Bushnik T, et al, 19(4):314–328

G

Gabapentin

Acute care management of post-TBI spasticity, Zafonte R, Elovic EP, Lombard L, 19(2):89–100

Gait

Ecological validity of walking speed assessment after traumatic brain injury: a pilot study, Moseley AM, Lanzarone S, Bosman JM, et al, 19(4):341–348

Evaluation and management of spastic gait in patients with traumatic brain injury, Esquenazi A, 19(2):109–118

Outcome assessment for spasticity management in the patient with traumatic brain injury: the state of the art, Elovic EP, Simone LK, Zafonte R, 19(2):155–177

Gamma aminobutyric acid (GABA)

Orally delivered baclofen to control spastic hypertonia in acquired brain injury, Meythaler JM, Clayton W, Davis LK, Guin-Renfroe S, Brunner RC, 19(2):101–108

Growth hormone deficiency

Anterior pituitary dysfunction after traumatic brain injury: Part II, Elovic EP, 19(2):184–187

H

Headache

Clinical caveats on medical assessment and treatment of pain after TBI, Ivanhoe CB, Hartman ET, 19(1):29–39

Occipital nerve blocks in postconcussive headaches: a retrospective review and report of ten patients, Hecht JS, 19(1):58–71

The problem of pain, Nicholson K, Martelli MF, 19(1):2–9

Psychological and neuropsychological integration in multidisciplinary pain management after TBI, Branca B, Lake AE III, 19(1):40–57

Head injury

Movement disorders after head injury, O'Suilleabhain P, Dewey RB, Jr., 19(4):305–313

Health support

Development of a telephone follow-up program for individuals following traumatic brain injury, Bell KR, Hoffman JM, Doctor JN, et al, 19(6):502–512

Heterotopic ossification

Clinical caveats on medical assessment and treatment of pain after TBI, Ivanhoe CB, Hartman ET, 19(1):29–39

Neuro-orthopedic management of shoulder deformity and dysfunction in brain-injured patients: A novel approach, Keenan MAE, Mehta S, 19(2):143–154

Hypernasality

An evaluation of continuous positive airway pressure (CPAP) therapy in the treatment of hypernasality following traumatic brain injury: a report of 3 cases, Cahill LM, Turner AB, Stabler PA, et al, 19(3):241–253

Hypothyroidism

Anterior pituitary dysfunction after traumatic brain injury: Part II, Elovic EP, 19(2):184–187

I

Imidazolines

Acute care management of post-TBI spasticity, Zafonte R, Elovic EP, Lombard L, 19(2):89–100

Interdisciplinary education

Collaboration between cognitive science and cognitive rehabilitation: a call for action, Powell JM, Hunt E, Pepping M, 19(3):266–276

Internet

An assistive technology intervention for verbose speech after traumatic brain injury: a single case study, Kirsch NL, Shenton M, Spirl E, et al, 19(5):366–377

L

Laryngeal function

Clinical characteristics of acute dysphagia in pediatric patients following traumatic brain injury, Morgan A, Ward E, Murdoch B, 19(3):226–240

Legislative issues

The Annual Congressional Brain Injury Awareness Fair, Bérubé JE, 19(1):82–84

M

Malingering

Determination of effort level, exaggeration, and malingering in neurocognitive assessment, Lynch WJ, 19(3):277–283

Medicaid

Fighting the silent epidemic: the Florida Brain and Spinal Cord Injury Program, Stuart M, 19(4):329–340

Memory

Change and predictors of change in communication, cognition, and social function between 1 and 5 years after traumatic brain injury, Hammond FM, Hart T, Bushnik T, et al, 19(4):314–328

Memory aids

Portable electronic devices as memory and organizational aids after traumatic brain injury: a consumer survey study, Hart T, Buchhofer R, Vaccaro M, 19(5):351–365

Model systems

Severity of injury and service utilization following traumatic brain injury: the first 3 months, Phillips VL, Greenspan AI, Stringer AY, Stroble AK, Lehtonen S, 19(3):217–225

Mood disorders

Psychological and neuropsychological integration in multidisciplinary pain management after TBI, Branca B, Lake AE III, 19(1):40–57

Motor point block

Clinical caveats on medical assessment and treatment of pain after TBI, Ivanhoe CB, Hartman ET, 19(1):29–39

Movement disorders

Movement disorders after head injury, O'Suilleabhain P, Dewey RB, Jr., 19(4):305–313

Myoclonus

Movement disorders after head injury, O'Suilleabhain P, Dewey RB, Jr., 19(4):305–313

N

Needs assessment

Perceived needs following traumatic brain injury, Corrigan JD, Whiteneck G, Mellick D, 19(3):205–216

Neurocognitive assessment

Determination of effort level, exaggeration, and malingering in neurocognitive assessment, Lynch WJ, 19(3):277–283

Neuropathic pain

Clinical caveats on medical assessment and treatment of pain after TBI, Ivanhoe CB, Hartman ET, 19(1):29–39

Neuropsychology

Neuropsychological performance and sleep disturbance following traumatic brain injury, Mahmood O, Rapport LJ, Hanks RA, Fichtenberg NL, 19(5):378–390

A new neuropsychological test battery: the NAB, Lynch WJ, 19(2):180–183

Psychological and neuropsychological integration in multidisciplinary pain management after TBI, Branca B, Lake AE III, 19(1):40–57

O

Occipital nerve block

Occipital nerve blocks in postconcussive headaches: a retrospective review and report of ten patients, Hecht JS, 19(1):58–71

Occupational issues

Perceived needs following traumatic brain injury, Corrigan JD, Whiteneck G, Mellick D, 19(3):205–216

Oral-motor skills

Clinical characteristics of acute dysphagia in pediatric patients following traumatic brain injury, Morgan A, Ward E, Murdoch B, 19(3):226–240

Outcomes

Change and predictors of change in communication, cognition, and social function between 1 and 5 years after traumatic brain injury, Hammond FM, Hart T, Bushnik T, et al, 19(4):314–328

Identifying environmental factors that influence the outcomes of people with traumatic brain injury, Whiteneck GG, Gerhart KA, Cusick CP, 19(3):191–204

Medical and social issues related to posttraumatic seizures in persons with traumatic brain injury, Bushnik T, Englander J, Duong T, 19(4):296–304

Participation as an outcome of traumatic brain injury rehabilitation, Cicerone KD, 19(6):494–501

Participation objective, participation subjective: a measure of participation combining outsider and insider perspectives, Brown M, Dijkers PJM, Gordon WA, et al, 19(6):459–481

Outcomes measurement

Latent factors in measures of rehabilitation outcomes after traumatic brain injury, Corrigan JD, Bogner J, 19(6):445–458

Outcome assessment for spasticity management in the patient with traumatic brain injury: the state of the art, Elovic EP, Simone LK, Zafonte R, 19(2):155–177

P

Pain/pain management

Clinical caveats on medical assessment and treatment of pain after TBI, Ivanhoe CB, Hartman ET, 19(1):29–39

Occipital nerve blocks in postconcussive headaches: a retrospective review and report of ten patients, Hecht JS, 19(1):58–71

Pain pathoetiology After TBI: Neural and nonneural mechanisms, Walker WC, 19(1):72–81

The problem of pain, Nicholson K, Martelli MF, 19(1):2–9

Psychological, neuropsychological, and medical considerations in assessment and management of pain, Martelli MF, Zasler ND, Bender MC, Nicholson K, 19(1):10–28

Psychological and neuropsychological integration in multidisciplinary pain management after TBI, Branca B, Lake AE III, 19(1):40–57

Parkinsonism

Movement disorders after head injury, O'Suilleabhain P, Dewey RB, Jr., 19(4):305–313

Participation

Participation as an outcome of traumatic brain injury rehabilitation, Cicerone KD, 19(6):494–501

Participation objective, participation subjective: a measure of participation combining outsider and insider perspectives, Brown M, Dijkers PJM, Gordon WA, et al, 19(6):459–481

Toward an explanatory model of social participation for adults with traumatic brain injury, Dumont C, Gervais M, Fougeyrollas P, Bertrand R, 19(6):431–444

Pediatric brain injury

Clinical characteristics of acute dysphagia in pediatric patients following traumatic brain injury, Morgan A, Ward E, Murdoch B, 19(3):226–240

Medical costs of mild to moderate traumatic brain injury in children, Brener I, Harman JS, Kelleher KJ, Yeates KO, 19(5):405–412

Visuomotor response time in children with a mild traumatic brain injury, Gagnon I, Swaine B, Friedman D, Forget R, 19(5):391–404

Personal digital assistant (PDA)

An assistive technology intervention for verbose speech after traumatic brain injury: a single case study, Kirsch NL, Shenton M, Spirl E, et al, 19(5):366–377

Personality disorders

Psychological and neuropsychological integration in multidisciplinary pain management after TBI, Branca B, Lake AE III, 19(1):40–57

Pharmacology update

Anterior pituitary dysfunction after traumatic brain injury: Part II, Elovic EP, 19(2):184–187

Benzodiazepines and antipsychotics: cognitive side effects, Perna R, 19(6):516–518

Physical therapy

Ecological validity of walking speed assessment after traumatic brain injury: a pilot study, Moseley AM, Lanzarone S, Bosman JM, et al, 19(4):341–348

Pituitary dysfunction, anterior

Anterior pituitary dysfunction after traumatic brain injury: Part II, Elovic EP, 19(2):184–187

Post-concussion syndrome

Occipital nerve blocks in postconcussive headaches: a retrospective review and report of ten patients, Hecht JS, 19(1):58–71

Psychological and neuropsychological integration in multidisciplinary pain management after TBI, Branca B, Lake AE III, 19(1):40–57

Posttraumatic epilepsy

Evaluation of seizure-like episodes in survivors of moderate and severe traumatic brain injury, Hudak AM, Trivedi K, Harper CR, et al, 19(4):290–295

Medical and social issues related to posttraumatic seizures in persons with traumatic brain injury, Bushnik T, Englander J, Duong T, 19(4):296–304

Posttraumatic headache

The problem of pain, Nicholson K, Martelli MF, 19(1):2–9

Psychological and neuropsychological integration in multidisciplinary pain management after TBI, Branca B, Lake AE III, 19(1):40–57

Problem solving

Change and predictors of change in communication, cognition, and social function between 1 and 5 years after traumatic brain injury, Hammond FM, Hart T, Bushnik T, et al, 19(4):314–328

Psychology

Collaboration between cognitive science and cognitive rehabilitation: a call for action, Powell JM, Hunt E, Pepping M, 19(3):266–276

Psychotherapy

Psychological and neuropsychological integration in multidisciplinary pain management after TBI, Branca B, Lake AE III, 19(1):40–57

Q

Quality of life

Outcome assessment for spasticity management in the patient with traumatic brain injury: the state of the art, Elovic EP, Simone LK, Zafonte R, 19(2):155–177

Participation as an outcome of traumatic brain injury rehabilitation, Cicerone KD, 19(6):494–501

Questionnaires

Characteristics of facility-based community integration programs for people with brain injury, Glenn MR, Goldstein R, Selleck EA, Rotman M, 19(6):482–493

Participation objective, participation subjective: a measure of participation combining outsider and insider perspectives, Brown M, Dijkers PJM, Gordon WA, et al, 19(6):459–481

R

Rehabilitation

An assistive technology intervention for verbose speech after traumatic brain injury: a single case study, Kirsch NL, Shenton M, Spirl E, et al, 19(5):366–377

Characteristics of facility-based community integration programs for people with brain injury, Glenn MR, Goldstein R, Selleck EA, Rotman M, 19(6):482–493

Collaboration between cognitive science and cognitive rehabilitation: a call for action, Powell JM, Hunt E, Pepping M, 19(3):266–276

Development of a telephone follow-up program for individuals following traumatic brain injury, Bell KR, Hoffman JM, Doctor JN, et al, 19(6):502–512

Fighting the silent epidemic: the Florida Brain and Spinal Cord Injury Program, Stuart M, 19(4):329–340

Latent factors in measures of rehabilitation outcomes after traumatic brain injury, Corrigan JD, Bogner J, 19(6):445–458

Outcome assessment for spasticity management in the patient with traumatic brain injury: the state of the art, Elovic EP, Simone LK, Zafonte R, 19(2):155–177

Participation as an outcome of traumatic brain injury rehabilitation, Cicerone KD, 19(6):494–501

Participation objective, participation subjective: a measure of participation combining outsider and insider perspectives, Brown M, Dijkers PJM, Gordon WA, et al, 19(6):459–481

Rehabilitative management of patients with disorders of consciousness: grand rounds, Giacino JT, Trott CT, 19(3):254–265

Severity of injury and service utilization following traumatic brain injury: the first 3 months, Phillips VL, Greenspan AI, Stringer AY, Stroble AK, Lehtonen S, 19(3):217–225

Research

Collaboration between cognitive science and cognitive rehabilitation: a call for action, Powell JM, Hunt E, Pepping M, 19(3):266–276

Ecological validity of walking speed assessment after traumatic brain injury: a pilot study, Moseley AM, Lanzarone S, Bosman JM, et al, 19(4):341–348

Research on minimally conscious patients: innovation or exploitation? Winslade WJ, 19(2):178–179

Research with brain-injured subjects, Tovino SA, 19(6):513–515

Resiliency factors

Toward an explanatory model of social participation for adults with traumatic brain injury, Dumont C, Gervais M, Fougeyrollas P, Bertrand R, 19(6):431–444

S

Science

Collaboration between cognitive science and cognitive rehabilitation: a call for action, Powell JM, Hunt E, Pepping M, 19(3):266–276

Secondary hypoadrenalism

Anterior pituitary dysfunction after traumatic brain injury: Part II, Elovic EP, 19(2):184–187

Seizures

Evaluation of seizure-like episodes in survivors of moderate and severe traumatic brain injury, Hudak AM, Trivedi K, Harper CR, et al, 19(4):290–295

Medical and social issues related to posttraumatic seizures in persons with traumatic brain injury, Bushnik T, Englander J, Duong T, 19(4):296–304

Self-efficacy

Toward an explanatory model of social participation for adults with traumatic brain injury, Dumont C, Gervais M, Fougeyrollas P, Bertrand R, 19(6):431–444

Service utilization

Severity of injury and service utilization following traumatic brain injury: the first 3 months, Phillips VL, Greenspan AI, Stringer AY, Stroble AK, Lehtonen S, 19(3):217–225

Sex education

An evaluation of sex education and information resources and their provision to adults with traumatic brain injury, Simpson G, Long E, 19(5):413–428

Shoulder disorders

Neuro-orthopedic management of shoulder deformity and dysfunction in brain-injured patients: A novel approach, Keenan MAE, Mehta S, 19(2):143–154

Shoulder pain

Clinical caveats on medical assessment and treatment of pain after TBI, Ivanhoe CB, Hartman ET, 19(1):29–39

Sleep disturbance

Neuropsychological performance and sleep disturbance following traumatic brain injury, Mahmood O, Rapport LJ, Hanks RA, Fichtenberg NL, 19(5):378–390

Social function

Change and predictors of change in communication, cognition, and social function between 1 and 5 years after traumatic brain injury, Hammond FM, Hart T, Bushnik T, et al, 19(4):314–328

Social issues

Medical and social issues related to posttraumatic seizures in persons with traumatic brain injury, Bushnik T, Englander J, Duong T, 19(4):296–304

Social participation

Participation as an outcome of traumatic brain injury rehabilitation, Cicerone KD, 19(6):494–501

Toward an explanatory model of social participation for adults with traumatic brain injury, Dumont C, Gervais M, Fougeyrollas P, Bertrand R, 19(6):431–444

Spasticity

Acute care management of post-TBI spasticity, Zafonte R, Elovic EP, Lombard L, 19(2):89–100

Clinical caveats on medical assessment and treatment of pain after TBI, Ivanhoe CB, Hartman ET, 19(1):29–39

Evaluation and management of spastic gait in patients with traumatic brain injury, Esquenazi A, 19(2):109–118

Neuro-orthopedic management of shoulder deformity and dysfunction in brain-injured patients: a novel approach, Keenan MAE, Mehta S, 19(2):143–154

Orally delivered baclofen to control spastic hypertonia in acquired brain injury, Meythaler JM, Clayton W, Davis LK, Guin-Renfroe S, Brunner RC, 19(2):101–108

Outcome assessment for spasticity management in the patient with traumatic brain injury: the state of the art, Elovic EP, Simone LK, Zafonte R, 19(2):155–177

Speech

An assistive technology intervention for verbose speech after traumatic brain injury: a single case study, Kirsch NL, Shenton M, Spirl E, et al, 19(5):366–377

Stroke

Orally delivered baclofen to control spastic hypertonia in acquired brain injury, Meythaler JM, Clayton W, Davis LK, Guin-Renfroe S, Brunner RC, 19(2):101–108

Subluxation

Neuro-orthopedic management of shoulder deformity and dysfunction in brain-injured patients: a novel approach, Keenan MAE, Mehta S, 19(2):143–154

T

Telehealth

Development of a telephone follow-up program for individuals following traumatic brain injury, Bell KR, Hoffman JM, Doctor JN, et al, 19(6):502–512

Tests

Determination of effort level, exaggeration, and malingering in neurocognitive assessment, Lynch WJ, 19(3):277–283

A new neuropsychological test battery: the NAB, Lynch WJ, 19(2):180–183

Tics

Movement disorders after head injury, O'Suilleabhain P, Dewey RB, Jr., 19(4):305–313

Traumatic brain injury. *See* Brain injury

Tremors

Movement disorders after head injury, O'Suilleabhain P, Dewey RB, Jr., 19(4):305–313

U

Upper motoneuron syndrome

Choosing upper limb muscles for focal intervention after traumatic brain injury, Mayer NH, 19(2):119–142

Evaluation and management of spastic gait in patients with traumatic brain injury, Esquenazi A, 19(2):109–118

V

Vegetative state

Rehabilitative management of patients with disorders of consciousness: grand rounds, Giacino JT, Trott CT, 19(3):254–265

Velopharyngeal dysfunction (VPD)

An evaluation of continuous positive airway pressure (CPAP) therapy in the treatment of hypernasality following traumatic brain injury: a report of 3 cases, Cahill LM, Turner AB, Stabler PA, et al, 19(3):241–253

Video-EEG monitoring

Evaluation of seizure-like episodes in survivors of moderate and severe traumatic brain injury, Hudak AM, Trivedi K, Harper CR, et al, 19(4):290–295

Visuomotor response time

Visuomotor response time in children with a mild traumatic brain injury, Gagnon I, Swaine B, Friedman D, Forget R, 19(5):391–404

W

Walking

Ecological validity of walking speed assessment after traumatic brain injury: a pilot study, Moseley AM, Lanzarone S, Bosman JM, et al, 19(4):341–348

Whiplash

Occipital nerve blocks in postconcussive headaches: A retrospective review and report of ten patients, Hecht JS, 19(1):58–71

Will (determination)

Toward an explanatory model of social participation for adults with traumatic brain injury, Dumont C, Gervais M, Fougeyrollas P, Bertrand R, 19(6):431–444